

Logic

Discrete Mathematics

Number Theory

Topic 01 — Logic

Mathematical Proofs

Lecture 01 — Introduction to Propositional Logic

Dr Kieran Murphy 

Recurrence Relations

Department of Computing and Mathematics,
Waterford IT.
(kmurphy@wit.ie)

Set Theory

Autumn Semester, 2021

Outline

- Propositions and fundamental logical operators (AND, OR and NOT).
- Evaluating logical expression using truth tables.
- Satisfiability, Tautologies and Contradictions.

Thought for the day ...

While walking through a fictional forest, you encounter three identical trolls guarding a bridge. Each troll is either a knight, who always tells the truth, or a knave, who always lies. The trolls will not let you pass until you correctly identify each as either a knight or a knave. Each troll makes a single statement:

If I am a knave, then there are exactly two knights here.



Troll 1 is lying.



Either we are all knaves or at least one of us is a knight..



Which troll are knights? and which are knaves?

1. Introduction 3
 - Propositional logic is concerned with analysing propositions (true or false statements).
 - A proposition may be atomic or compound (build up using logical connectives).
 - Constructing compound propositions using *And*, *Or* and *Not*.
2. Truth tables 13
 - Evaluating an expression for all possible input combinations.
3. Tautologies and Contradictions 21
 - Statements that are always true or always false.

Logic

Logic is “science of reasoning”

- Allows us to represent knowledge in precise, unambiguous way.
- Allows us to make valid inferences using a set of consistent rules.
- Roots of logic date back to the ancient Greeks, e.g., Aristotle.
- Greeks were interested in valid logical inference rules, such as syllogisms:

“All men are mortal.

Socrates is a man.

Therefore, Socrates is mortal.”



The Partially Examined Life podcast: www.partiallyexaminedlife.com

The Fallacy-a-Day Podcast: <http://fallacyaday.com>

Propositional Logic

- The building blocks of propositional logic are propositions

Definition 1 (Proposition)

A **proposition** (**statement**) is a sentence that is either **True** or **False**.

- Examples:

“Java is a programming language.”

True

“Cork is the capital of Ireland.”

False

“ $1 + 2 = 3$ ”

True

“Today is Tuesday.”

depends

“The universe is fine-tuned.”

unknown (at present)

- Examples of sentences that are not propositions/statements:

- “How are you?”* — A question cannot be assign a **True/False** value.
- “Stop sleeping in class!”* — An order cannot be assign a **True/False** value.
- “Correct horse battery staple.”* — Not a sentence.
- “This sentence is false.”* — Pathological example.

Propositional Variables, Truth Value

Given a proposition we are interested in knowing its **truth value**.

Definition 2 (Truth Value)

The **truth value** of a proposition identifies whether a proposition is true (written **True** or **T** or 1) or false (written **False** or **F** or 0)

Question

What is truth value of “*Tuesday in the day after Sunday*” ?

F

Notation

- Variables that represent propositions are called propositional variables.
- Denote propositional variables using lower-case letters, such as $p, p_1, p_2, q, r, s, \dots$
- Truth value of a propositional variable is either **T** or **F**.

Compound vs Atomic Propositions

- Propositional logic allows constructing more complex propositions from atomic ones.
- More complex propositions formed using **logical connectives** (also called **boolean connectives** or **logical operators**).
- The three basic logical connectives:

Connective	Symbol	Python
conjunction (AND)	\wedge	and
disjunction (OR)	\vee	or
negation (NOT)	\neg	not

- Propositions formed using these logical connectives called **compound propositions**; otherwise called **atomic propositions**.

Today is wet and I am hungry

Compound vs Atomic Propositions

- Propositional logic allows constructing more complex propositions from atomic ones.
- More complex propositions formed using **logical connectives** (also called **boolean connectives** or **logical operators**).
- The three basic logical connectives:

Connective	Symbol	Python
conjunction (AND)	\wedge	and
disjunction (OR)	\vee	or
negation (NOT)	\neg	not

- Propositions formed using these logical connectives called **compound propositions**; otherwise called **atomic propositions**.

Today is wet and I am hungry

Exercise

Classify each of the sentences below as an atomic statement, a compound statement, or not a statement at all.

- 1 The sum of the first 100 odd positive integers.
- 2 Everybody needs somebody sometime.
- 3 Waterford will win the All-Ireland or I'll eat my hat.
- 4 Go to your room!
- 5 Every natural number greater than 1 is either prime or composite.
- 6 This sentence is false.

Exercise

Classify each of the sentences below as an atomic statement, a compound statement, or not a statement at all.

- ① The sum of the first 100 odd positive integers.

—*This is not even a sentence (no verb).*

- ② Everybody needs somebody sometime.

—*This is an atomic statement.*

- ③ Waterford will win the All-Ireland or I'll eat my hat.

—*This is a compound statement.*

- ④ Go to your room!

—*This is an order, not a statement*

- ⑤ Every natural number greater than 1 is either prime or composite.

—*This is a compound statement.*

- ⑥ This sentence is false.

—*This is sentence but is not a statement.*

Negation (NOT)

- **Negation** of a proposition, p , written, $\neg p$, represents the proposition:
“It is not the case that p .”
- What is the relationship between the truth value of p and $\neg p$?

If p is **T**, then $\neg p$ is **F** and vice versa.

- In simple English, what is $\neg p$ if p stands for ...

p	$\neg p$
<hr style="width: 100%;"/> <i>“Today is Tuesday.”</i> <i>“1 + 1 = 2”</i>	<hr style="width: 100%;"/> <i>“Today is not Tuesday.”</i> <i>“1 + 1 \neq 2”</i>

- Properties of NOT
 - $\neg \neg p = p$

Conjunction (AND)

- **Conjunction** of two propositions, p and q , written as $p \wedge q$, is the proposition:

“p and q”

- What is the relationship between the truth value of p and of q and the truth value of $p \wedge q$?

$$p \wedge q = \begin{cases} \mathbf{T} & \text{if both } p \text{ is } \mathbf{T} \text{ and } q \text{ is } \mathbf{T} \\ \mathbf{F} & \text{otherwise} \end{cases}$$

Example

What is the conjunction and the truth value of $p \wedge q$ for ...

- $p =$ “It is a autumn semester”, $q =$ “Today is Thursday”
- $p =$ “It is Tuesday”, $q =$ “It is morning”

Disjunction (OR)

- **Disjunction** of two propositions, p and q , written as $p \vee q$, is the proposition

“ p or q ”

- What is the relationship between the truth value of p and of q and the truth value of $p \vee q$?

$$p \vee q = \begin{cases} \mathbf{T} & \text{if either } p \text{ is } \mathbf{T} \text{ or } q \text{ is } \mathbf{T}, \text{ or both are } \mathbf{T} \\ \mathbf{F} & \text{otherwise} \end{cases}$$

Example

What is the disjunction and the truth value of $p \vee q$ for ...

- $p =$ “*It is a autumn semester*”, $q =$ “*Today is Thursday*”
- $p =$ “*It is Friday*”, $q =$ “*It is morning*”

Python supports the fundamental logical connectives (programmers call them “logical operators”)

Logical Connective	Math	Python Operator
conjunction (AND)	\wedge	and
disjunction (OR)	\vee	or
negation (NOT)	\neg	not

1. Introduction 3
 - Propositional logic is concerned with analysing propositions (true or false statements).
 - A proposition may be atomic or compound (build up using logical connectives).
 - Constructing compound propositions using *And*, *Or* and *Not*.
2. Truth tables 13
 - Evaluating an expression for all possible input combinations.
3. Tautologies and Contradictions 21
 - Statements that are always true or always false.

Propositional Formulas and Truth Tables

- A **propositional formula** is logical expression constructed from atomic and compound propositions and logical connectives.
- A **truth table** for a propositional formula, A , shows the truth value of A for every possible value of its constituent atomic propositions.

Negation		Conjunction			Disjunction		
p	$\neg p$	p	q	$p \wedge q$	p	q	$p \vee q$
F	T	F	F	F	F	F	F
F	T	F	T	F	F	T	T
T	F	T	F	F	T	F	T
T	F	T	T	T	T	T	T



NOT



AND



OR

Truth tables and Logic Gates

AND

NOT

OR

p	$\neg p$

p	q	$p \wedge q$


p	q	$p \vee q$

Truth tables and Logic Gates

AND

NOT

p	$\neg p$
F	T



OR

p	q	$p \wedge q$


p	q	$p \vee q$

Truth tables and Logic Gates

AND

NOT

p	$\neg p$
F	T
T	F




OR

p	q	$p \wedge q$

p	q	$p \vee q$


Truth tables and Logic Gates

AND




NOT

p	$\neg p$
F	T
T	F



OR





p	q	$p \wedge q$
F	F	F

p	q	$p \vee q$
F	F	F



Truth tables and Logic Gates

AND





NOT

p	$\neg p$
F	T
T	F






OR





p	q	$p \wedge q$
F	F	F
F	T	F

p	q	$p \vee q$
F	F	F
F	T	T

Truth tables and Logic Gates




AND


p	$\neg p$
F	T
T	F

NOT






p	q	$p \wedge q$
F	F	F
F	T	F
T	F	F

p	q	$p \vee q$
F	F	F
F	T	T
T	F	T

OR




Truth tables and Logic Gates

AND



p	$\neg p$
F	T
T	F





NOT



p	q	$p \wedge q$
F	F	F
F	T	F
T	F	F
T	T	T

p	q	$p \vee q$
F	F	F
F	T	T
T	F	T
T	T	T


OR




Other Resources

iPad/iPhone Apps (assume similar on Android)

Circuit Coder



Boolean Master



Videos

- <https://class.coursera.org/cs101/lecture/17>
Part of the Computer Science 101 by Nick Parlante on coursera.

Constructing Truth Tables

Useful strategy for constructing truth tables for a formula:

- STEP 1 Identify the constituent atomic propositions of the formula.
- STEP 2 Identify compound propositions in within the formula in increasing order of complexity, including the formula itself.
- STEP 3 Construct a table enumerating all combinations of truth values for atomic propositions.
- STEP 4 Fill in values of compound propositions for each row.

Examples

Construct truth tables for the following formulas:

1 $(p \vee q) \wedge \neg p$

2 $(p \wedge q) \vee (\neg p \wedge \neg q)$

3 $(p \vee q \vee \neg r) \wedge r$

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p q

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$p \vee q$	$\neg p$	$(p \vee q) \wedge \neg p$

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$p \vee q$	$\neg p$	$(p \vee q) \wedge \neg p$
F	F			
F	T			
T	F			
T	T			

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$p \vee q$	$\neg p$	$(p \vee q) \wedge \neg p$
F	F	F	T	F
F	T			
T	F			
T	T			

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$p \vee q$	$\neg p$	$(p \vee q) \wedge \neg p$
F	F	F	T	F
F	T	T	T	T
T	F			
T	T			

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$p \vee q$	$\neg p$	$(p \vee q) \wedge \neg p$
F	F	F	T	F
F	T	T	T	T
T	F	T	F	F
T	T			

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$p \vee q$	$\neg p$	$(p \vee q) \wedge \neg p$
F	F	F	T	F
F	T	T	T	T
T	F	T	F	F
T	T	T	F	F

Example 1: $(p \vee q) \wedge \neg p$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$p \vee q$	$\neg p$	$(p \vee q) \wedge \neg p$
F	F	F	T	F
F	T	T	T	T
T	F	T	F	F
T	T	T	F	F

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p q

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$(p \wedge q)$	$\neg p$	$\neg q$	$(\neg p \wedge \neg q)$	$(p \wedge q) \vee (\neg p \wedge \neg q)$

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$(p \wedge q)$	$\neg p$	$\neg q$	$(\neg p \wedge \neg q)$	$(p \wedge q) \vee (\neg p \wedge \neg q)$
F	F					
F	T					
T	F					
T	T					

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$(p \wedge q)$	$\neg p$	$\neg q$	$(\neg p \wedge \neg q)$	$(p \wedge q) \vee (\neg p \wedge \neg q)$
F	F	F	T	T	T	T
F	T					
T	F					
T	T					

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$(p \wedge q)$	$\neg p$	$\neg q$	$(\neg p \wedge \neg q)$	$(p \wedge q) \vee (\neg p \wedge \neg q)$
F	F	F	T	T	T	T
F	T	F	T	F	F	F
T	F					
T	T					

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$(p \wedge q)$	$\neg p$	$\neg q$	$(\neg p \wedge \neg q)$	$(p \wedge q) \vee (\neg p \wedge \neg q)$
F	F	F	T	T	T	T
F	T	F	T	F	F	F
T	F	F	F	T	F	F
T	T					

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$(p \wedge q)$	$\neg p$	$\neg q$	$(\neg p \wedge \neg q)$	$(p \wedge q) \vee (\neg p \wedge \neg q)$
F	F	F	T	T	T	T
F	T	F	T	F	F	F
T	F	F	F	T	F	F
T	T	T	F	F	F	T

Example 2: $(p \wedge q) \vee (\neg p \wedge \neg q)$

STEP 1 Identify the constituent atomic propositions ... p and q

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	$(p \wedge q)$	$\neg p$	$\neg q$	$(\neg p \wedge \neg q)$	$(p \wedge q) \vee (\neg p \wedge \neg q)$
F	F	F	T	T	T	T
F	T	F	T	F	F	F
T	F	F	F	T	F	F
T	T	T	F	F	F	T

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

Example 3: $(p \vee q \vee \neg r) \wedge r$

STEP 1 Identify the constituent atomic propositions ... p , q , and r

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p q r

Example 3: $(p \vee q \vee \neg r) \wedge r$

STEP 1 Identify the constituent atomic propositions ... p , q , and r

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$

Example 3: $(p \vee q \vee \neg r) \wedge r$

STEP 1 Identify the constituent atomic propositions ... p , q , and r

STEP 2 Identify compound propositions ...

STEP 3 Enumerate all combinations of truth values for atomic propositions ...

STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F			
F	F	T			
F	T	F			
F	T	T			
T	F	F			
T	F	T			
T	T	F			
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T			
F	T	F			
F	T	T			
T	F	F			
T	F	T			
T	T	F			
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F			
F	T	T			
T	F	F			
T	F	T			
T	T	F			
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F	T	T	F
F	T	T			
T	F	F			
T	F	T			
T	T	F			
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F	T	T	F
F	T	T	F	T	T
T	F	F			
T	F	T			
T	T	F			
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F	T	T	F
F	T	T	F	T	T
T	F	F	T	T	F
T	F	T			
T	T	F			
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F	T	T	F
F	T	T	F	T	T
T	F	F	T	T	F
T	F	T	F	T	T
T	T	F			
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1 Identify the constituent atomic propositions ... p , q , and r
- STEP 2 Identify compound propositions ...
- STEP 3 Enumerate all combinations of truth values for atomic propositions ...
- STEP 4 Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F	T	T	F
F	T	T	F	T	T
T	F	F	T	T	F
T	F	T	F	T	T
T	T	F	T	T	F
T	T	T			

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1** Identify the constituent atomic propositions ... p , q , and r
- STEP 2** Identify compound propositions ...
- STEP 3** Enumerate all combinations of truth values for atomic propositions ...
- STEP 4** Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F	T	T	F
F	T	T	F	T	T
T	F	F	T	T	F
T	F	T	F	T	T
T	T	F	T	T	F
T	T	T	F	T	T

Example 3: $(p \vee q \vee \neg r) \wedge r$

- STEP 1** Identify the constituent atomic propositions ... p , q , and r
- STEP 2** Identify compound propositions ...
- STEP 3** Enumerate all combinations of truth values for atomic propositions ...
- STEP 4** Fill in values of compound propositions for each row ...

p	q	r	$\neg r$	$(p \vee q \vee \neg r)$	$(p \vee q \vee \neg r) \wedge r$
F	F	F	T	T	F
F	F	T	F	F	F
F	T	F	T	T	F
F	T	T	F	T	T
T	F	F	T	T	F
T	F	T	F	T	T
T	T	F	T	T	F
T	T	T	F	T	T

1. Introduction 3
 - Propositional logic is concerned with analysing propositions (true or false statements).
 - A proposition may be atomic or compound (build up using logical connectives).
 - Constructing compound propositions using *And*, *Or* and *Not*.
2. Truth tables 13
 - Evaluating an expression for all possible input combinations.
3. Tautologies and Contradictions 21
 - Statements that are always true or always false.

Introduction to Propositional Logic — Summary

1. Introduction 3
 - Propositional logic is concerned with analysing propositions (true or false statements).
 - A proposition may be atomic or compound (build up using logical connectives).
 - Constructing compound propositions using *And*, *Or* and *Not*.
2. Truth tables 13
 - Evaluating an expression for all possible input combinations.
3. Tautologies and Contradictions 21
 - Statements that are always true or always false.

Satisfiable, Tautologies and Contradictions

Satisfiable

A proposition is **satisfiable** if it is **True** for at least one set of inputs (case).

Tautology

A **tautology** is an expression involving logical variables that is **True** in all cases.

- Examples

- $p \vee \neg p$

“Tomorrow, I will be dead or I will be alive”

- $(p \wedge q) \vee (p \wedge \neg q) \vee \neg p$

Contradiction

A **contradiction** is an expression involving logical variables that is **False** in all cases.

- Examples

- $p \wedge \neg p$

“On Friday, I will win the lottery and not win the lottery.”