

Logic

# Discrete Mathematics

Number Theory

Mathematical Proofs

Topic 06 — Graphs and Networks

Lecture 01 — Fundamental Concepts in Graph Theory

Dr Kieran Murphy   

Recurrence Relations

Department of Computing and Mathematics,  
Waterford IT.  
([kmurphy@wit.ie](mailto:kmurphy@wit.ie))

Set Theory

Autumn Semester, 2021

## Outline

- Fundamental graph concepts and definitions
- A selection of common graphs

Enumeration

# Outline

---

1. Graph Theory at a Glance	2
1.1 History of Graph Theory	3
1.2 What is a Graph?	6
1.3 Applications	7
2. Graph Jargon	8
3. Fundamental Concepts	22
4. Some Common Graphs	32




# History of Graph Theory — Stage 1: Birth (circa 1735)

## The Bridges of Königsberg Problem

Is it possible to walk through the city of Königsberg that would cross each of the seven bridges once and only once and return to one's starting point.

## Solved by Euler

- Lead to the birth of graph theory.



# History of Graph Theory — Stage 2: Mathematical Interest

Problems were only of mathematical interest, for example


## Cycles in Polyhedra



Thomas P. Kirkman



William R. Hamilton




Hamiltonian cycles in Platonic graphs


# History of Graph Theory — Stage 3: Applications

## In Physics ...

### Trees in Electric Circuits



Gustav Kirchhoff



## In Chemistry ...

### Enumeration of Chemical Isomers





Arthur Cayley



James J. Sylvester




George Polya




## In Computing ..., in Biology ..., in Social Science ..., in ...



### Protein-Protein Interaction Network



### Scientific collaboration Network




### Friendship Network



# Graphs — An Informal Definition

A **graph** is a set of objects with pairwise connections.

- The notion of a **graph** is deceptively simple: It is a collection of points (called **vertices** or **nodes**) that are joined by lines (called **edges** or **arcs**).
- All that matters about an edge is which two vertices it connects — and sometimes its length, capacity and/or cost — but not the layout of the edge.



# Graph Applications

Graph theory is one of the most widely applicable areas of mathematics.

<b>Application</b>	<b>Vertices</b>	<b>Edges</b>
communication	telephones, computers	cables, connections
circuits	gates, registers, processors	wires
mechanical	joints	rods, beams, springs
financial	stocks, currency	transactions
games	board positions	legal moves
transportation	road intersections, airports	roads, air routes
scheduling	tasks	precedence constraints
software systems	functions	function calls
social relationship	people, actors	friendships, movie casts
chemical compounds	molecules	bonds
hydraulic	reservoirs, pumping stations	pipelines
Internet	web pages	hyperlinks
neural networks	neurons	synapses

# Outline

---

1. Graph Theory at a Glance	2
2. Graph Jargon	8
2.1 Common Terms	12
2.2 Typical Graph Problems	20
3. Fundamental Concepts	22
4. Some Common Graphs	32




# Graph — Formal Definition

## Definition 1 (Simple Graph)


A **simple graph**,  $G$ , consists of a non-empty finite set,  $V(G)$ , of elements called **vertices** (or **nodes**), and a finite set,  $E(G)$ , of distinct unordered pairs of distinct elements of  $V(G)$  called **edges** (or **arcs**).

- The set  $V(G)$  is called the **vertex set** and has  $n = |V(G)|$  elements (vertices).
- The set  $E(G)$  the **edge set** of  $G$ . and has  $m = |E(G)|$  elements (edges).
- An edge  $(v, w)$  is said to **join** the vertices  $v$  and  $w$ , and is usually abbreviated to  $vw$  or  $v-w$ .
- The simple graph on the right has vertex set  $V(G) = \{u, v, w, z\}$  and edge set  $E(G) = \{uv, uw, vw, wz\}$ .



# General Graph

In any simple graph there is at most one edge joining any given pair of vertices (no **multiple edges**), and all edges join distinct vertices (no **loops**). There are situations where these restrictions are not desirable. (e.g., adding redundancy in a communication network.)



## Definition 2 (General Graph)

A **general graph**,  $G$ , consists of a non-empty finite set,  $V(G)$ , of elements called **vertices**, and a finite multi-set (or family),  $E(G)$ , of unordered, not necessarily distinct, elements of  $V(G)$  called **edges**.

- In the theory component of this course we will try to prove results for general graphs but we will be more restrictive in our Python implementations and mainly concentrate on simple graphs.

# Warning — No Standard Notation

Graph notation is not standard!

Some of the different variations that are popular are:

- **Wilson:1996**

- **simple graph** — A graph with no multiple edges and no loops.
- **general graph** or **graph** — A graph with multiple edges and/or loops.

- **Sedgwick:2003**

- **graph** — A graph with no multiple edges and no loops.
- **multi-graph** — A graph with multiple edges and/or loops.


- **Trudeau:1993**

- **graph** — A graph with no multiple edges and no loops.
- **multi-graph** — A graph with multiple edges.
- **pseudo-graph** — A graph with multiple edges and loops.

# Graph Jargon — Vertex

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components




This graph has 13 vertices:

$\{A, B, C, D, E, F, G, H, I, J, K, L, M\}$

# Graph Jargon — Edge

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components




This graph has 13 edges:

$$\{(A, B), (A, C), (A, G), (A, F), (D, E), (D, F), (E, F), (E, G), (H, I), (J, K), (J, L), (J, M), (L, M)\}$$

# Graph Jargon — Parallel Edges, Self Loop

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components




This graph has two parallel edges  $\{(A, F), (A, F)\}$  and two self-loops  $\{(H, H), (G, G)\}$ .

# Graph Jargon — Directed Graph

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components




The edges in a directed graph (**digraph**) are drawn with an arrow indicating the direction. So for example, this graph has edge  $(A, B)$  but not edge  $(B, A)$ .

# Graph Jargon — Weighted Graph

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components




The edges in a weighted graph have weights representing length, cost, or delay in traversing that edge.



# Graph Jargon — Path, Cycle

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components




A **path** is a sequence of connected vertices, e.g.,  $\{D, E, F, A, G\}$ .

A **cycle** is a path with same end vertices, e.g.,  $\{J, L, M, J\}$ .

# Graph Jargon — Tree, Forest

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components




A **tree** is graph with no cycles.

A **forest** is a set of trees.

# Graph Jargon — Connected Components

## Terminology:

- Vertex
- Edge
- Parallel edges, self loop
- Directed graph
- Weighted graph
- Path, cycle
- Tree, forest
- Connected, connected components



A graph is **connected** if there is a path between any two vertices.

Parts of a graph that are not connected to each other are called **connected components**.

# Typical Graph Problems

---

## Paths

**Path:** Is there a path between two nodes,  $u$  and  $v$ ?

**Shortest Path:** What is the shortest path between  $u$  and  $v$ ?

**Longest Path:** What is the longest path between  $u$  and  $v$ ?

## Cycles and Tours

**Cycle:** Is there a cycle in the graph?

**Euler Tour:** Is there a cycle path that uses each edge exactly once?

**Hamilton Tour:** Is there a cycle path that uses each vertex exactly once?

## Connectivity

**Connectivity:** Is it possible to connect all of the vertices?

**MST:** What is the optimum way to connect all of the vertices?

**Bi-connectivity:** Is there a vertex whose removal disconnects the graph?

# Typical Graph Problems



## Planarity

**Planarity:** Is it possible to draw the graph in the plane with no crossing edges?

**Depth:** What is the minimum number of crossing need to layout a non-planar graph.

## Graph Colouring

**Vertex Colouring:** What is the minimum number of colours needed to colour the graph vertices so that adjacent vertices have different colours?

**Edge Colouring:** What is the minimum number of colours needed to colour the graph edges so that adjacent edges have different colours?

# Outline

---

1. Graph Theory at a Glance	2
2. Graph Jargon	8
<b>3. Fundamental Concepts</b>	<b>22</b>
3.1 Isomorphic Graphs	23
3.2 Labelled and Unlabelled Graphs	25
3.3 Connectedness and Adjacency	26
3.4 Degree and Related Concepts	29
4. Some Common Graphs	32

# Isomorphism

Given two graphs are they equal\* or distinct?

— Remember a graph is not changed by rearranging its drawing layout.

## Definition 3 (Isomorphism)

Two graphs  $G_1$  and  $G_2$  are **isomorphic** if there is a one-to-one correspondence between the vertices of  $G_1$  and  $G_2$  such that the number of edges joining any two vertices of  $G_1$  is equal to the number of edges joining the corresponding vertices of  $G_2$ .

- For many problems, the labels on the vertices are unnecessary and can be disregarded. In this case we say that the two **unlabelled graphs** are isomorphic if we can assign labels to both graphs so that the resulting **labelled graphs** are isomorphic.



---

\*Isomorphic = Greek “equal” + “shape”

# Example — Isomorphic Graphs

## Example 4

The two graphs shown bellow are isomorphic under the correspondence  $u \leftrightarrow 5, v \leftrightarrow 3, w \leftrightarrow 1, x \leftrightarrow 4, y \leftrightarrow 2, z \leftrightarrow 6$ .




It is easier to check this by building the one-to-one correspondence a step at a time:  $u \leftrightarrow 5, v \leftrightarrow 3, w \leftrightarrow 1, x \leftrightarrow 4, y \leftrightarrow 2, z \leftrightarrow 6$ .




# Labelled and Unlabelled Graphs

The number of permutations of vertex labels grows large rapidly as the number of vertices in the graph increases. For example, for simple graphs with three vertices, there are only four distinct unlabelled simple graphs while there are eight distinct labelled simple graphs.



Unlabelled simple graphs



Labelled simple graphs

This difference increases as the number of vertices increase, for example, consider graphs with four vertices, here there are 6 distinct unlabelled simple graphs and 64 distinct labelled simple graphs with four vertices, (see Wilson, page 10).

# Connectness

Graphs can be combined to make a larger graph.

## Definition 5 (Graph union)

If the two graphs are  $G_1 = (V(G_1), E(G_1))$  and  $G_2 = (V(G_2), E(G_2))$ , where  $V(G_1)$  and  $V(G_2)$  are disjoint, then their **union**,  $G_1 \cup G_2$  is the graph with vertex set  $V(G_1) \cup V(G_2)$  and edge multi-set  $E(G_1) \cup E(G_2)$ .


## Definition 6 (Connected)

A graph is **connected** if it cannot be expressed as the union of two graphs, and **disconnected** otherwise.

- Any disconnected graph,  $G$ , can be expressed as the union of connected graphs, each of which is a **component** of  $G$ .

# Example

The graph,  $G$ , depicted bellow has three components



Hence we have  $G = G_1 \cup G_2 \cup G_3$  where

$$G_1 = (\{A, B, C, D, E, F, G\}, \{A-B, A-C, A-F, A-G, D-E, D-F, E-F, E-G\})$$

$$G_2 = (\{H, I\}, \{H-I\})$$



$$G_3 = (\{J, K, L, M\}, \{J-K, J-L, J-M, L-M\})$$

# Adjacency

## Definition 7 (Adjacent/Incident)

Two vertices,  $v$  and  $w$ , are **adjacent** if there is an edge,  $v-w$ , joining them, and the vertices  $v$  and  $w$  are then **incident** with such an edge.

Similarly, two distinct edges,  $e$  and  $f$ , are **adjacent** if they have a vertex in common.



# Degree and Related Concepts

## Definition 8 (Degree)


The **degree** of a vertex,  $v$ , is the number of edges incident with  $v$ , denoted by  $\deg(v)$ .

- Normal convention is that a loop at  $v$  contributes 2 (rather than 1) to the degree of  $v$ .
- A vertex of degree 0 is an **isolated vertex**, and a vertex of degree 1 is an **end-vertex**.
- A graph in which all vertices have degree  $r$  is said to be **regular of degree  $r$** .
- The degree sequence of a graph,  $G$ , consists of degree of all of the vertices of  $G$  sorted in non-decreasing order.

# Example

## Example 9

Consider the following graph



- Vertices  $B$ ,  $C$ ,  $H$ ,  $I$ , and  $K$  are **end-vertices**.
- Vertex  $A$  has degree 4.
- The degree sequence of the graph is  $1, 1, 1, 1, 1, 2, 2, 2, 2, 3, 3, 3, 4$ .

# Review Exercises 1 (Fundamental Concepts)

## Question 1:

Draw a graph with degree sequence  $(3, 3, 5, 5, 5, 5)$ . Does there exist a *simple* graph with this degree sequence? Justify your answer.

## Question 2:

State, with an explanation, which of the following sequences are the degree sequences of a simple graph. For those sequence that are degree sequences, draw a simple graph with that degree sequence.

(a)  $(1, 1, 2, 2, 3, 4, 4, 5, 5,)$

(b)  $(2, 2, 2, 3, 5, 6, 6, 6)$

(c)  $(1, 2, 3, 4, 5, 6, 7)$

## Question 3:

If  $G$  is a simple graph with at least two vertices, prove that  $G$  must contain two or more vertices of the same degree.

## Question 4:

**(Hard)** Is it possible for two *different* (non-isomorphic) graphs to have the same number of vertices and the same number of edges? What if the degrees of the vertices in the two graphs are the same (so both graphs have vertices with degrees 1, 2, 2, 3, and 4, for example)? Draw two such graphs or explain why not.

## Question 5:


Are the two graphs below equal? Are they isomorphic? If they are isomorphic, give the isomorphism. If not, explain.

Graph 1:

$$V = \{a, b, c, d, e\}$$

$$E = \{\{a, b\}, \{a, c\}, \{a, e\}, \{b, d\}, \{b, e\}, \{c, d\}\}.$$

Graph 2:



# Outline

---

1. Graph Theory at a Glance	2
2. Graph Jargon	8
3. Fundamental Concepts	22
<b>4. Some Common Graphs</b>	<b>32</b>
4.1 Null Graphs	33
4.2 Complete Graphs	34
4.3 Cycle, Wheel and Path Graphs	35
4.4 Regular Graphs	36
4.5 Bipartite Graphs	37
4.6 Other Graphs	39




# Null Graphs

## Definition 10 (Null graph, $N_n$ )

A graph whose edge-set is empty is a **null graph**. A null graph with  $n$  vertices is denoted by  $N_n$ .  $(n \geq 1)$

- The vertices of a null graph are all isolated.

## Example 11 (The null graph, $N_4$ )





# Complete Graphs

## Definition 12 (Complete graph, $K_n$ )

A simple graph in which every pair of vertices are adjacent is a **complete graph**. A complete graph with  $n$  vertices is denoted by  $K_n$ .  $(n \geq 1)$

- A complete graph of  $n$  vertices has  $n(n - 1)/2$  edges.

## Example 13 (The complete graphs, $K_4$ and $K_5$ )





# Cycle Graphs

## Definition 14 (Cycle graph, $C_n$ )

A connected graph that is regular of degree 2 is a **cycle graph**. A cycle graph with  $n$  vertices is denoted by  $C_n$ .  $(n \geq 3)$

- The graph obtain from  $C_n$  by removing an edge is the **path graph** on  $n$  vertices, denoted by  $P_n$ .  $(n \geq 3)$
- The graph obtain from  $C_{n-1}$  by joining each vertex to a new vertex  $v$  is called the **wheel graph** on  $n$  vertices, denoted by  $W_n$ .  $(n \geq 4)$

## Example 15 (Graphs $C_6$ , $P_6$ and $W_6$ )




# Regular Graphs

## Definition 16 (Regular graph)

A graph in which all of the vertices are of degree  $r$  is a **regular graph of degree  $r$**  or  **$r$ -regular**.

- A **cubic graph** is a regular of degree 3 graph.
  - A important example of a cubic graph is the **Petersen graph**.
- The null graph,  $N_n$ , is a regular graph of degree 0.
- The complete graph,  $K_n$ , is a regular graph of degree  $n - 1$ .

## Example 17 (The Petersen graph)






# Bipartite Graphs

## Definition 18 (Bipartite graph)

A graph,  $G$ , in which its vertex set can be split into two disjoint sets,  $A$  and  $B$ , so that each edge of  $G$  joins a vertex of  $A$  and a vertex of  $B$  is a **bipartite graph**.

- An alternative definition: A **bipartite graph** is a graph whose vertices can be coloured black and white so that each edge joins a black vertex and a white vertex.
- A **complete bipartite graph** is a bipartite graph in which each vertex in  $A$  is joined to each vertex in  $B$  by just one edge, i.e., all black vertices are joined to all white vertices, and all white vertices are joined to all black vertices.
- A complete bipartite graph with  $r$  black vertices and  $s$  white vertices is denoted by  $K_{r,s}$ .

# Examples of Bipartite Graphs



# Other Graphs


- Gallery of named graphs

[http://en.wikipedia.org/wiki/Gallery\\_of\\_named\\_graphs](http://en.wikipedia.org/wiki/Gallery_of_named_graphs)

- ISEM's MATtours:

[http:](http://www.hamline.edu/~lcofes/SciMathMN/gallery.html)

[//www.hamline.edu/~lcofes/SciMathMN/gallery.html](http://www.hamline.edu/~lcofes/SciMathMN/gallery.html)




Balaban 10-cage

Balaban 11-cage

Bidiakis cube

Brinkmann graph

Bull graph




Butterfly graph

Chvátal graph

Diamond graph

Dürer graph

Ellingham–Horton 54-graph




Ellingham–Horton 78-graph

Errera graph

Franklin graph

Frucht graph

Goldner–Harary graph



# Review Exercises 2 (Some Common Graphs)





## Question 1:

Give an example of, or explain why it doesn't exist, each of the following.

- (a) A bipartite graph that is regular of degree 5.
- (b) A complete graph that is a wheel.
- (c) A cubic graph with 11 vertices.
- (d) A graph (other than  $K_5$ ,  $K_{4,4}$ , or  $Q_4$ ) that is regular of degree 4.

## Question 2:

Which of the graphs below are bipartite? Justify your answers.



## Question 3:

For which  $n \geq 3$  is the graph  $C_n$  bipartite?